

Article type: Original Research Article

Title: The dietary flavonol quercetin ameliorates angiotensin II-induced redox signalling imbalance in a human umbilical vein endothelial cell model of endothelial dysfunction via ablation of p47^{phox} expression.

Authors: Huw S. Jones^{1,2}, Andrew Gordon², Simba G. Magwenzi², Khalid Naseem², Stephen L. Atkin^{2,3}, Fraser L. Courts^{2,4}

Affiliations: ¹Department of Sports, Health and Exercise Sciences, University of Hull, Hull, HU6 7RX, UK; ²Centre for Cardiovascular and Metabolic Research, Hull York Medical School, University of Hull, Hull, HU6 7RX, UK, ³Weill Cornell Medical College Qatar, Doha, Qatar; ⁴Campden BRI, Station Road, Chipping Campden, Gloucestershire, GL55 6LD, UK.

Corresponding author: Dr Huw S. Jones

Department of Sports, Health and Exercise Sciences, University of Hull, Hull, East Riding of Yorkshire, UK, HU6 7RX

Telephone number: +44 (0) 1482

Fax number: +44 (0) 1482

huw.jones@hull.ac.uk

Key words (max 5): Quercetin, nitric oxide, superoxide, endothelial cells, NADPH oxidase

22 **Abstract**

23 Scope: Quercetin is reported to reduce blood pressure in hypertensive but not normotensive
 24 humans, but the role of endothelial redox signalling in this phenomenon has not been
 25 assessed. This study investigated the effects of physiologically-obtainable quercetin
 26 concentrations in a human primary cell model of endothelial dysfunction in order to elucidate
 27 the mechanism of action of its antihypertensive effects.

28 Methods and results: Angiotensin II (100 nM, 8 h) induced dysfunction, characterised by
 29 suppressed nitric oxide availability ($85 \pm 4\%$ $p < 0.05$) and increased superoxide production
 30 ($136 \pm 5\%$, $p < 0.001$). These effects were ablated by an NADPH oxidase inhibitor. Quercetin
 31 ($3 \mu\text{M}$, 8 h) prevented angiotensin II induced changes in nitric oxide and superoxide levels,
 32 but no effect on upon nitric oxide or superoxide in control cells. The NADPH oxidase
 33 subunit p47^{phox} was increased at the mRNA and protein levels in angiotensin II-treated cells
 34 ($130 \pm 14\%$ of control, $p < 0.05$), which was ablated by quercetin co-treatment. Protein kinase
 35 C activity was increased after angiotensin II treatment ($136 \pm 51\%$), however this was
 36 unaffected by quercetin co-treatment.

37 Conclusions: Physiologically-obtainable quercetin concentrations are capable of ameliorating
 38 angiotensin II-induced endothelial nitric oxide and superoxide imbalance via protein kinase
 39 C-independent restoration of p47^{phox} gene and protein expression.

40

41

42

43

44 **Introduction**

45 Evidence from a wide variety of studies suggests that particular dietary polyphenolic plant
46 secondary metabolites are capable of modulating clinical markers of cardiovascular health in
47 humans (1-5). Of the many thousands of these structures present in fruit and vegetable-rich
48 diets, the flavonol quercetin is amongst the most ubiquitous, being found most commonly in
49 berries, apples and onions, and being freely available as an isolate marketed as a nutritional
50 supplement. Quercetin has been shown to reduce blood pressure in hypertensive but not
51 normotensive human volunteers (6-10). These studies showed significant decreases of both
52 systolic and diastolic blood pressure in hypertensive individuals, apparently independent of
53 brachial artery vasodilation, angiotensin converting enzyme activity, and plasma nitrite
54 concentrations (7, 8). However, the molecular mechanism underpinning the potential beneficial
55 effects of quercetin have remained elusive. In models of rodent hypertension modulation of
56 NADPH oxidase expression and reactive oxygen species (ROS) formation (11, 12) have been
57 suggested as key targets for quercetin. However, most of these experiments have used rodent
58 aortic ring segments which do not allow specific assessment of endothelial function (13-15).

59 The endothelium is an important source of vasodilatory and constrictive signalling factors
60 involved in the downstream regulation of blood pressure and normal vascular function.
61 Notably, the redox signalling agents nitric oxide (NO^\cdot) and superoxide ($\text{O}_2^{\cdot-}$) rapidly and
62 antagonistically mediate such effects (16). Imbalance in this redox equilibrium is strongly
63 implicated in the pathophysiology of several cardiovascular diseases, and it may therefore be
64 hypothesised that restoration of endothelial function plays a significant role in mediating the
65 effects of quercetin in endothelial dysfunction. Indeed, relaxation studies using stimulated
66 aortic ring segments have suggested the involvement of endothelial NO^\cdot availability in the
67 mechanism of action of quercetin, and a cell culture model of HOCl -induced endothelial
68 dysfunction has shown that supraphysiological concentrations of quercetin can increase NOS

activity (13-15, 17). Interestingly, this suggestion is in apparent conflict with the later study of Larson *et al.* (2012), which concluded that the clinical effects of quercetin in hypertensive individuals were independent of plasma nitrite concentrations; a clinical marker of circulating NO[•] availability. In addition, the barrier function of the endothelium makes this structure most likely to be exposed to plasma quercetin following oral administration.

This study therefore aimed to determine the potential effects of physiologically-obtainable quercetin concentrations on NO[•] and O₂^{•-} signalling imbalance. For this study primary vascular endothelial cells (Human Umbilical Vein Endothelial Cells; HUVEC) have been used rather than endothelial cell lines, which have been reported to vary in characteristics quite dramatically when compared to *ex vivo* tissues. HUVEC are well characterised, relatively easily accessible, primary cells that have been reported to exhibit key endothelial cell functions, and are extensively used in the study of the effects of dietary chemicals upon cardiovascular biology and diseases (18-24). Specifically, we have assessed the effects of quercetin on NO[•] availability, O₂^{•-} production and NADPH oxidase subunit protein levels and activity. The concentrations of quercetin added have been chosen to replicate those which may arise from dietary intake, providing meaningful information on the mechanisms which underlie clinical observations. This use of physiologically-obtainable quercetin concentrations is an important part of the novelty of this study as although the role of NADPH oxidase and NO[•] production in the mechanism of action has been previously investigated in a variety of model systems, these studies have mainly assessed supra-physiological concentrations of quercetin. Thus there is a possibility that the previously reported effects of quercetin upon vascular dysfunction are actually due to off-target effects and are therefore not relevant to the mechanism of action in humans. Our study aims to address this important consideration.

93 **Materials and methods**

94 *Materials*

95 Unless otherwise listed below, all reagents were sourced from Sigma-Aldrich (Poole, UK).

96 Human Umbilical Vein Endothelial Cells were purchased from Promocell (Heidelberg,
97 Germany) as cryopreserved aliquots of 5×10^5 cells pooled from 4 donors. Endothelial Cell
98 Growth Medium (ECGM) was also purchased from Promocell (Heidelberg, Germany).
99 Medium 199 (M199) was obtained from Life Technologies (Paisley, UK). Nunc Nunclon
100 plasma-treated black 96-well plates were bought from Fisher Scientific (Leicestershire, UK).
101 4, 5-Diaminofluorescein diacetate (DAF-2 DA) was purchased from Enzo Life Sciences
102 (Exeter, UK). The DC protein assay kit, 2x Lamelli buffer, Mini-Protean 12% TGX gels, 10x
103 TGS buffer, 0.2 μ m PVDF Turboblot membrane packs, Precision Plus protein marker ladder,
104 Clarity Western ECL solution, Aurum total RNA extraction kit and iScript cDNA synthesis kit
105 were all obtained from Bio-Rad (Hertfordshire, UK). Marvel low-fat powdered milk was
106 purchased from a local supermarket. Autoradiography films (Amersham ECL) were purchased
107 from Fisher Scientific (Loughborough, UK). An Endothelin-1 ELISA assay kit was purchased
108 from Enzo Life Sciences (Exeter, UK). Solaris PCR expression assays for NOS3 (eNOS),
109 NCF-1, EF1 α and β -actin, and the Solaris qPCR mastermix were purchased from GE
110 Healthcare (Buckinghamshire, UK).

111 The following antibodies were used: p47^{phox} (Abcam, Cambridge, UK, Ab63361, Lot #
112 803556), gp91^{phox} (Abcam, Cambridge, UK, AB129068, Lot # GR83718-3), β -actin (Abcam,
113 Cambridge, UK, ab20272, Lot # GR88824-1), Phospho-(Ser) PKC substrate antibody (Cell
114 Signalling, 2261, Lot # 18), anti-beta tubulin (Millipore, 06-661, Lot # 239882), Goat anti-
115 rabbit HRP-conjugate (Sigma-Aldrich, Poole, UK, A6154, Lot # 090116176), Goat anti-mouse

HRP-conjugate (Abcam, Cambridge, UK, ab97023 , Lot # GR87150-8), Goat anti-rabbit HRP-conjugate (GE Healthcare, RPN4301, Lot # 9526414).

Human Umbilical Vein Endothelial Cell (HUVEC) culture

HUVEC were grown to ~95% confluence with ECGM + 20% foetal bovine serum (FBS) over 3-4 d, with the culture media refreshed every 2 d. After trypsinisation, HUVEC were seeded into either 96-well or 6-well plates at a density of 14,700 cells/cm² and cultured for the durations described below and in the figure legends, with media refreshed every 2 days.

Live cell measurement of NO[•] availability in HUVEC cultures

Passage 4 – 6 HUVEC were seeded in 96-well plates (5 x 10³ cells / well) and allowed to proliferate for 2 - 3 d as required. Cell cultures were serum-deprived for 24 hours in Medium-199 containing 0.5% FBS (M199) before endpoint measurement. Ang II was prepared as a 1 mM stock in ddH₂O (18.2 MΩ), diluted in culture media to produce 10 nM – 10 μM final concentration. 8 hr before end-point measurement cells were treated in M199 with or without Ang II. Cells were treated with quercetin or solvent only as directed in the Figure legends for 8 hours in the presence or absence of Ang II. For cells treated with L-NAME, a stock solution of 100 mM was prepared in DMSO, and diluted 1000-fold in culture media. Cells were treated with L-NAME, or a vehicle control, for 30 minutes before end-point measurement. After incubation with the appropriate treatment(s), cells were washed once on a warmed plate with 200 μl of warmed HBSS (containing magnesium and calcium), and 70 μl of warm HBSS (containing magnesium and calcium) containing 2 μM DAF-2DA was added per well. For L-NAME treated cells, L-NAME was also included in this dye solution. Fluorescence (λ_{ex} 485

nm, λ_{em} 520 nm) was measured every 3 min over a 30 minute period using a Tecan infinite X200 plate reader, maintained at 37°C, with a matrix of 4x4 points per well measured (ten flashes per point), and a manual gain of 100 set. Linear rates ($r^2 \geq 0.99$) were calculated for all wells and a mean rate of fluorescence was calculated for each treatment (n=4-6 wells per treatment). Wells containing cells only, and wells without cells were also assessed

Protein extraction from HUVEC cultures

Cells were cultured as described and treated 8 h before extraction with either DMF (0.1% v/v), quercetin (3 μ M), DMF (0.1% v/v) and Ang II (100 nM), or quercetin (3 μ M) and Ang II (100 nM), in M199 containing 0.5% FBS. Monolayers were washed twice in 3 ml of PBS per well, before the addition of 200 μ l of RIPA buffer (1% NP-40 substitute, 0.5% w/v sodium deoxycholate, 0.1% w/v sodium dodecyl sulphate, made up in PBS containing protease inhibitor cocktail) per well. Cells were harvested by scraping before incubation for 5 min on ice to aid lysis. The cell lysates were sonicated (Sonic Vibracell VCX130PB) for 3 x 10 s bursts on ice. Lysates were centrifuged at 16 100 x g, 5 min, and the resulting supernatant stored at -20°C.

Western blotting for NADPH oxidase subunits $p47^{phox}$ and $gp91^{phox}$

Protein extracts, prepared as above, were quantified using the DC protein assay as described by the manufacturer. 20 μ g of protein extract was added in a 1:1 ratio to 2x Lamelli buffer. The resulting samples were then boiled for 5 minutes at 95°C, before incubation on ice for 2 minutes, and centrifugation at 16 100 x g for 30 seconds. Proteins were separated using 12% TGX gel for 30 minutes at 250 V in 1x TGS buffer. Separated proteins were transferred PVDF membranes before washing in 1x TBST buffer (20 mM Trizma base, 137 mM sodium chloride,

1 ml of tween20, pH 7.6) and blocking in a solution of 5% w/v milk in 1x TBST for 30 minutes. Membranes were rinsed and washed 3x 10 minutes in 1x TBST buffer, before incubation for 1 hour with the appropriate primary antibody dilution (p47^{phox} – 1:2000; gp91^{phox} – 1:5000) in 1% w/v milk in 1x TBST buffer. Following further rinsing and 3 x 10 minute washes in 1x TBST buffer, membranes were incubated for 1 hour with the appropriate secondary antibody (Goat anti-rabbit HRP conjugate – 1:30000) diluted in 1% w/v milk in 1x TBST. The membranes were then rinsed and washed again (3x 10 minutes in 1x TBST) and processed by ECL. Membranes were stored at 4°C for use in measuring β -actin levels as a loading control. Membranes were incubated for 2x 10 minute washes in stripping buffer (200 mM glycine, 3.5 mM sodium dodecyl sulphate, 1% tween20, adjusted to pH 2.2), washed twice in TBST, and then blocked (5% w/v milk, 30 minutes) and washed a further 3 times. The membranes were incubated with the β -actin HRP conjugated antibody (1:2 000 000 dilution) in 1% w/v milk in 1xTBST buffer for 1 hour before a further 3x 10 minutes washes in 1x TBST buffer. The membranes were incubated in ECL solution and developed as previously described, with a 2.5 minute exposure to the autoradiography film.

For the assessment of PKC-phosphorylated substrate consensus sequences membranes were blocked using 10% bovine serum albumin (BSA) for 1 h followed by incubation with a PKC-phosphorylated substrate consensus sequence primary antibody diluted to 1:1000 in 1 % BSA for 1 h. An appropriate HRP-conjugated secondary antibody diluted 1:10,000 in 1% BSA was used to detect the primary antibody. Membranes were stripped and re-probed with for β -tubulin at a dilution of 1:1000 overnight at 4°C.

All films were digitised and densitometry was done using the ImageJ software. A ratio of protein of interest: β -actin was calculated for each sample with the control samples for each biological replicate set at 100%. For the PKC-phosphorylated substrate films, the entire lane was quantified by densitometry as described above, and normalised to β -tubulin.

187 *Intracellular O₂⁻ determination by LC-MS detection of 2HE⁺*

188 HUVEC were cultured as described in 6-well plates and serum-deprived in M199 containing
189 0.5% FBS for 24 h prior to experiments. At 8 h before experiments these cultures were
190 incubated with or without Ang II (100 nM) in the presence of vehicle (0.1% v/v DMF), 3 nM,
191 or 3 μ M quercetin, with 3 wells per treatment. For experiments using the NADPH oxidase
192 inhibitor VAS-2870, cells were treated with or without Ang II for 8 h, with either vehicle (0.1%
193 DMF) or 1 μ M VAS-2870 for 30 minutes before assay. After 8 h incubation the cells were
194 washed twice in excess warm PBS containing 100 μ M DTPA and then incubated in darkness
195 with 2 ml of 20 μ M dihydroethidium (DHE, dissolved in DMSO at a stock concentration of 20
196 mM) in HBSS for 30 min. Following two further washes with excess PBS - DTPA, each well
197 was scraped in 500 μ l of methanol, and wells from the same treatment groups were pooled and
198 stored overnight at -20 °C. The resulting lysates were centrifuged (16,100 x g, 10 minutes, 4
199 °C) with both the supernatant and pellet retained separately. The supernatant was evaporated
200 to dryness using a centrifugal evaporator without heating (Genevac MiVac, Genevac, Ipswich,
201 UK), and the resulting residue was dissolved in 1:1 water:stabilisation solution (33.3%
202 methanol, 10 μ M fluorescein internal standard, 0.1 mM DTPA, and 5 mM sodium ascorbate,
203 made up to final volume with water) with a final volume of 120 μ l for LC-MS analysis. The
204 pellet was dissolved in 100 μ l of 0.1 M sodium hydroxide and the protein content was
205 quantified using the DC assay as directed by the manufacturer.

206 For LC-MS analysis of DHE oxidation products, the specific O₂⁻ reaction product (2-HE⁺)
207 was detected at a *m/z* ratio of 330.3, and at a retention time of 7 min. Separation was achieved
208 using a Shimadzu prominence LC20 quaternary pump and autosampler, a Shimadzu CTO10
209 column oven, and an Agilent Eclipse XDB-C18 column (5 μ m, 4.6 x 150 mm) at a flow rate
210 of 0.5 ml/min. Water containing 0.5% formic acid (solvent A) and methanol containing 0.5%
211 formic acid (solvent B) were used to achieve separation as follows; 43% B for 3.35 minutes,

increasing to 85% B by 8 minutes, maintained at 85% B until 11.5 minutes, reduced to 43% B by 11.75 minutes, held at 43% B until 18.75 minutes. Column oven temperature was set to 40 °C. The injection volume used was 20 µl. A Shimadzu LC2020 single quadrupole mass spectrometer was used as the detector with the following settings: Fluorescein was used as an internal standard, at a m/z ratio of 333.0 and a retention time of 11.25 minutes. Peaks corresponding to 2-HE⁺ and fluorescein were integrated and peak areas were calculated. The ratio of 2-HE⁺:fluorescein was normalised to the total protein. Control samples from each independent experiment were set to 100%, with all other treatments expressed as a percentage of control, and 3 independent experiments were analysed. A representative chromatogram is shown in Figure S1.

RNA extraction, cDNA synthesis and qPCR of eNOS and β -actin

HUVEC cultures were grown as described above in 6-well plate format, with and without Ang II (8 h, 100 nM) and quercetin (3 µM, 8 h) treatments. Cells were washed and scraped in PBS, with wells from the same treatment group pooled together, centrifuged, and RNA was extracted using the Bio-Rad Aurum RNA extraction kit as directed by the manufacturer. Eluted RNA was quantified by spectrophotometry. Using the Bio-Rad iScript cDNA synthesis kit, as directed by the manufacturer, 500 ng of RNA was reverse transcribed, and the resulting cDNA was quantified by spectrophotometry. Relative quantification of gene expression, comparing control against treated samples, was achieved using the validated Solaris PCR expression assay probes for eNOS, p47^{phox}, Elongation Factor 1 α and β -actin, by loading 150 ng of cDNA template to the assay probes and Solaris PCR master mix as directed by the manufacturer. The following PCR conditions were used for these probe sets: 1 cycle of 95°C 15 min followed by 50 cycles of 95°C for 15 sec and 60°C for 1 min. The resulting Ct values were analysed by the

$\Delta\Delta\text{Ct}$ method, using β -actin or Elongation Factor 1 α as the reference gene. Three independent experiments were analysed, with mean fold change in eNOS or p47^{phox} expression in treated samples compared to control samples calculated, \pm 1 standard deviation.

Statistical analysis

All statistical analyses used Sigmaplot version 12.0, with comparisons between multiple groups done by ANOVA or ANOVA on ranks, as appropriate after testing for normality of distribution and homogeneity of variance. Post-hoc tests were done using Tukey post-hoc tests. Values less than $p = 0.05$ were considered statistically significant.

Results*Ang II inhibits the production of NO[•] in HUVEC*

In order to examine the effects of quercetin on endothelial dysfunction we first exposed HUVEC to a range of Ang II concentrations (10 nM – 10 μ M) for up to 24 h, under optimised culture conditions (as outlined in Figure S2), and the effect of Ang II treatment upon NO[•] availability was determined (Figure 1 A-D). It was observed that concentrations below 10 μ M Ang II resulted in a 10 – 20% decrease in the rate of nitric oxide production, measured in live HUVEC cultures by determining the rate of fluorescence accumulation using diaminofluorescein (DAF). The greatest and most consistent reduction in NO[•] was observed for an 8 h incubation at 100 nM angiotensin II (Figure 1B). We did not observe a dose-response for Ang II in these experiments over the concentration range tested, however the observation of a lack of effect at higher concentrations (1 μ M and above) is consistent with previous research, as is the observed time dependency of Ang II treatment (18). Based upon these experiments future Ang II treatments were at a concentration of 100 nM for 8 h; a treatment regime that resulted in a decrease in NO[•] availability of $15 \pm 4\%$ compared to untreated cells (Figure 1E, $p < 0.05$). To confirm that this effect was mediated in a receptor dependent manner the experiments were repeated in the presence of the angiotensin receptor type 1 antagonist irbesartan (10 μ M, 8 h). Under these conditions no change in NO production was observed (Figure 1E). It was also noted that DMSO (0.1% v/v, 8 h), a commonly-used carrier solvent, attenuated the effects of Ang II on NO[•] levels in this system (Figure S2). Dimethyl-formamide (DMF) did not attenuate the effects of Ang II on NO[•] and was therefore used for all future experiments.

Physiologically-obtainable quercetin concentrations prevents Ang II-induced decreases in nitric oxide production

Using the live-cell nitric oxide assay described above, we next assessed the effect of quercetin upon the rate of nitric oxide production in HUVEC cultures with and without Ang II treatment (8 h, 100 nM). An initial test concentration of 3 μ M quercetin was selected for these experiments in order to test a physiologically-obtainable concentration (7, 25, 26). Quercetin (3 μ M, 8 h) was shown to prevent Ang II-induced decreases in the rate of nitric oxide production but had no effect on control cells (Figure 2A). A dose response of quercetin (3 nM – 3 μ M) in Ang II treated cells showed that quercetin prevented Ang II-induced dysfunction at all concentrations tested (Figure 2B). We next assessed the effect of quercetin treatment (3 μ M, 8 h) on the expression of endothelial Nitric Oxide Synthase (eNOS) using qPCR, with no changes in gene expression observed for any treatment compared to untreated cells (Figure 2C).

Quercetin prevents Ang II-induced superoxide production by NADPH oxidases

As NADPH oxidase expression and reactive oxygen species (ROS) formation have been suggested as key targets for quercetin (11, 12) we next sought to determine the effects of both Ang II and quercetin treatments on superoxide production in HUVEC cultures. Superoxide was increased by $36 \pm 5\%$ compared to control cells, and this was prevented by quercetin co-treatment at both 3 μ M and 3 nM (Figure 3A). The prevention of Ang II-induced superoxide production by the NADPH oxidase inhibitor VAS-2870 (Figure 3B, 1 μ M, 8 h, (27)) suggests that this ROS-producing complex was a major source of Ang II-induced superoxide in these HUVEC cultures.

Quercetin prevents Ang II-induced p47^{phox} expression at the protein and mRNA levels

In order to explore the potential mechanism by which quercetin prevents Ang II-induced dysfunction we next assessed the effect of quercetin treatment on the expression of the NADPH oxidase subunits p47^{phox} and gp91^{phox}. Quercetin (3 μ M, 8 h) prevented Ang II-induced p47^{phox} protein levels (130 \pm 14% of untreated cells), but had no effect upon p47^{phox} protein levels in control cells (Figure 4A). This trend was also reflected at the mRNA level (Figure 4B). The expression of gp91^{phox} protein was unaffected by either Ang II or quercetin treatments (Figure 4C).

Ang II-induced PKC activity is not modulated by quercetin co-treatment

Protein Kinase C activity is a well-characterised component of the Ang II signalling cascade and known activator of NADPH oxidase activity. Treatment of HUVEC with Ang II caused a trend of increase in PKC activity (116 \pm 32%, $p < 0.05$), which was unaffected by co-incubation of cell with quercetin (Figure 5). Quercetin alone (3 μ M, 8 h) appeared to have no PKC activity (Figure 5). These data suggest the possibility that the observed effects of quercetin are independent of PKC activity.

Discussion

In this study we have utilised angiotensin II to induce endothelial dysfunction *in vitro*, as characterised by decreased availability of NO[•], increased intracellular O₂^{•-}, increased NADPH oxidase subunit expression and increased PKC activity. Using this model we have probed the impacts of the flavonoid quercetin, which has previously been reported to modulate the expression of the NADPH oxidase subunit p47^{phox}, O₂^{•-} (SOD-inhibitable lucigenin detection), and of NO[•] production (nitrate and nitrite levels) *in vitro* (28, 29). However, these previous

studies used supraphysiological concentrations of quercetin under non-dysfunctional conditions and their findings are in conflict with clinical research. Consequently we have sought to address the discrepancy between the *in vitro* and clinical data by using physiologically-attainable quercetin concentrations and assessed effects under both non-dysfunctional and dysfunctional conditions.

Our study has shown that physiologically-obtainable concentrations of quercetin restore endothelial function (availability of NO[•], intracellular O₂^{•-} and p47^{phox} expression) to control levels in dysfunctional cells (induced by Ang II) whereas there are no effects in non-dysfunctional cells. This is in concordance with the current clinical data for the effects of quercetin in healthy and diseased populations (7, 8). The mechanism of action of quercetin that underlies these effects is currently unknown, with mechanistic insights from clinical studies and rodent studies being conflicting. Although these rodent studies have implicated changes in NADPH oxidase activity and increased availability of NO[•] as part of the mechanism of action, due to the inherent multicellular and multi-tissue complexity of the aortic ring segments, they have mainly focussed upon the effects of quercetin on vascular smooth muscle rather than the endothelium in *ex vivo* aortic ring segments. The findings presented from our study demonstrate that quercetin restores endothelial function in Ang II-treated cells and suggests that the observed restoration of endothelial NO[•] and O₂^{•-} signalling via the restoration of NADPH oxidase expression (possibly independently of PKC activity) is part of the mechanism of action of quercetin (Figure 6). Interestingly, over the concentration range tested (3 nM – 3 µM), there was no observed dose response and a similar ablation of endothelial dysfunction by quercetin was seen at all tested concentrations. There are several possible explanations for this observation including that the concentrations assessed in this study were not in the linear phase of a classical s-shaped dose response, or that quercetin potentially has multiple different mechanisms of action which can result in a variety of different shaped dose

response curves (30). Ultimately, future research should focus upon investigating the dose-response characteristics of quercetin in the context of endothelial dysfunction, as well as investigating the role of other potential mechanisms of action, such as other kinase signalling cascades (e.g. MAPK) and the phosphorylation control of eNOS and p47^{phox}.

Quercetin has been reported in human plasma in the nanomolar to low micromolar range, after dietary intake and supplementation (7, 8, 25, 26). Human pharmacokinetic studies also have shown that quercetin undergoes substantial conjugative metabolism, with the formation of sulfate and glucuronide conjugates, whereas the aglycone form is not detected in human plasma (31). It should be noted that these circulating conjugated forms are not currently widely available, and that as a result we have used the aglycone form of quercetin in this study. This is not necessarily a major limitation as it has recently been suggested that the anti-hypertensive effects of quercetin glucuronides are dependent on deconjugation activity and the liberation of quercetin (32). HUVEC have also been reported to express β -glucuronidase at the mRNA level (33), suggesting that they also possess at least some deconjugative machinery. Taken together, these data suggest that the use of the aglycone form of quercetin in the endothelial cell model reported is a valid approach for assessing its effects on endothelial dysfunction.

In summary, physiologically-obtainable concentrations of the dietary flavonoid quercetin restored NO[•] availability, O₂^{•-} production and p47^{phox} expression to control levels, possibly in a PKC-independent manner, in a human primary cell model of endothelial dysfunction. There were no further improvements in NO[•] availability under control conditions. These results are consistent with clinical observations that quercetin reduces blood pressure in hypertensive, but not normotensive, individuals, and suggest a potential mechanism by which quercetin mediates this effect.

Author contributions

Experiments were designed by HSJ and FLC. Experiments were done by HSJ, FLC, AG and SM. KN provided reagents and resources for the PKC experiments. Funding was provided by SLA. All authors reviewed and approved the final manuscript.

Acknowledgements

This work was co-funded by the InnovateUK and Boots Pharmaceuticals (TSB 101137), and HEIF5 funding. The authors would like to thank Prof. Henry Leese for his constructive comments on the manuscript, and Drs Roger Sturmev and Sam Xu for access to equipment.

Author disclosure statement

For all authors: No competing financial interests exist.

395 **List of Abbreviations**

396 Ang II – Angiotensin II

397 cDNA – Copy deoxyribose nucleic acid

398 DAF-2 DA – Diaminofluorescein diacetate

399 ddH₂O – distilled water

400 DHE - Dihydroethidium

401 DMF – Dimethyl-formamide

402 DMSO – Dimethyl-sulfoxide

403 DTPA – Diethylene triamine penta-acetic acid

404 ECGM – Endothelial cell growth media

405 ECL – Enhanced luminol chemiluminescence

406 eNOS – Endothelial nitric oxide synthase

407 FBS – Fetal bovine serum

408 HBSS – Hank's balanced salt solution

409 HRP – Horseradish peroxidase

410 HUVEC – Human umbilical vein endothelial cells

411 kDa – Kilo Dalton

412 LC-MS – Liquid chromatography mass spectrometry

413 L-NAME – NG-Nitro-L-arginine methyl ester

- 414 M199 – Medium 199
- 415 NADPH – Nicotinamide adenine dinucleotide 2'-phosphate (reduced form)
- 416 NO[•] - Nitric oxide
- 417 O₂^{•-} - Superoxide
- 418 PBS – Phosphate buffered saline
- 419 PKC – Protein kinase C
- 420 PVDF – Polyvinylidene difluoride
- 421 qPCR – Quantitative polymerase chain reaction
- 422 RNA – Ribose nucleic acid
- 423 ROS – Reactive oxygen species
- 424 SOD – Superoxide dismutase
- 425
- 426
- 427
- 428
- 429
- 430
- 431
- 432

433 **References**

- 434 1. Heiss C, Finis D, Kleinbongard P, Hoffmann A, Rassaf T, Kelm M, et al. Sustained increase in
435 FMD after daily intake of high flavanol cocoa drink over 1 week. *Journal of Cardiovascular*
436 *Pharmacology*. 2007;49:74-80.
- 437 2. Heiss C, Kleinbongard P, Dejam A, Perre S, Schroeter H, Sies H, et al. Acute consumption of
438 flavanol-rich cocoa and the reversal of endothelial dysfunction in smokers. *Journal of the American*
439 *College of Cardiology*. 2005;46(7):1276-83.
- 440 3. Duffy SJ, Keaney JF, Holbrook M, Gokce N, Swerdloff PL, Frei B, et al. Short- and Long-Term
441 Black Tea Consumption Reverses Endothelial Dysfunction in Patients With Coronary Artery Disease.
442 *Circulation*. 2001;104(2):151-6.
- 443 4. Hooper L, Kroon PA, Rimm EB, Cohn JS, Harvey I, Le Cornu KA, et al. Flavonoids, flavonoid-rich
444 foods, and cardiovascular risk: a meta-analysis of randomized controlled trials. *American Journal of*
445 *Clinical Nutrition*. 2008;88:38-50.
- 446 5. Ried K, Sullivan TR, Fakler P, Frank OR, Stocks NP. Effect of cocoa on blood pressure. *Cochrane*
447 *Database of Systematic Reviews*. 2012(8).
- 448 6. Loke WM, Hodgson JM, Proudfoot JM, McKinley AJ, Puddey IB, Croft KD. Pure dietary
449 flavonoids quercetin and (-)epicatechin augment nitric oxide products and reduce endothelin-1 in
450 acutely healthy men. *The American Journal of Clinical Nutrition*. 2008;88:1018-25.
- 451 7. Larson A, Witman MA, Guo Y, Ives S, Richardson RS, Bruno RS, et al. Acute, quercetin-induced
452 reductions in blood pressure in hypertensive individuals are not secondary to lower plasma
453 angiotensin-converting enzyme activity or endothelin-1: nitric oxide. *Nutr Res*. 2012;32(8):557-64.
- 454 8. Edwards RL, Lyon T, Litwin SE, Rabovsky A, Symons JD. Quercetin reduces blood pressure in
455 hypertensive subjects. *The Journal of nutrition*. 2007;137:2405-11.
- 456 9. Pfeuffer M, Auinger A, Bley U, Kraus-Stojanowic I, Laue C, Winkler P, et al. Effect of quercetin
457 on traits of the metabolic syndrome, endothelial function and inflammation in men with different
458 APOE isoforms. *Nutrition, metabolism, and cardiovascular diseases : NMCD*. 2013;23(5):403-9.
- 459 10. Egert S, Bosy-Westphal A, Seiberl J, Kurbitz C, Settler U, Plachta-Danielzik S, et al. Quercetin
460 reduces systolic blood pressure and plasma oxidised low-density lipoprotein concentrations in
461 overweight subjects with a high-cardiovascular disease risk phenotype: a double-blinded, placebo-
462 controlled cross-over study. *The British journal of nutrition*. 2009;102(7):1065-74.
- 463 11. Larson AJ, Symons JD, Jalili T. Quercetin: A Treatment for Hypertension?—A Review of Efficacy
464 and Mechanisms. *Pharmaceuticals*. 2010;3(1):237-50.
- 465 12. Larson AJ, Symons JD, Jalili T. Therapeutic potential of quercetin to decrease blood pressure:
466 review of efficacy and mechanisms. *Adv Nutr*. 2012;3(1):39-46.
- 467 13. Sanchez M, Galisteo M, Vera R, Villar IC, Zarzuelo A, Tamagro J, et al. Quercetin downregulates
468 NADPH oxidase increases eNOS activity and prevents endothelial dysfunction in spontaneously
469 hypertensive rats. *Journal of hypertension*. 2006;24:75-84.
- 470 14. Sanchez M, Lodi F, Vera R, Villar IC, Cogolludo A, Jimenez R, et al. Quercetin and isorhamnetin
471 prevent endothelial dysfunction, superoxide production, and overexpression of p47phox induced by
472 angiotensin II in rat aorta. *The Journal of nutrition*. 2007;137:910-5.
- 473 15. Khoo NK, White CR, Pozzo-Miller L, Zhou F, Constance C, Inoue T, et al. Dietary flavonoid
474 quercetin stimulates vasorelaxation in aortic vessels. *Free radical biology & medicine*. 2010;49(3):339-
475 47.
- 476 16. Drummond GR, Sobey CG. Endothelial NADPH oxidases which NOX to target in vascular
477 disease. *Trends in Endocrinology and Metabolism*. 2014;25(9):452-63.
- 478 17. Shen Y, Croft KD, Hodgson JM, Kyle R, Lee IL, Wang Y, et al. Quercetin and its metabolites
479 improve vessel function by inducing eNOS activity via phosphorylation of AMPK. *Biochem Pharmacol*.
480 2012;84(8):1036-44.
- 481 18. Rueckschloss U. Dose-Dependent Regulation of NAD(P)H Oxidase Expression by Angiotensin
482 II in Human Endothelial Cells: Protective Effect of Angiotensin II Type 1 Receptor Blockade in Patients

- With Coronary Artery Disease. *Arteriosclerosis, thrombosis, and vascular biology*. 2002;22(11):1845-51.
19. Ahn HY, Kim CH, Ha TS. Epigallocatechin-3-gallate Regulates NADPH Oxidase Expression in Human Umbilical Vein Endothelial Cells. *The Korean journal of physiology & pharmacology : official journal of the Korean Physiological Society and the Korean Society of Pharmacology*. 2010;14(5):325-9.
 20. Chae YJ, Kim CH, Ha TS, Heschler J, Ahn HY, Sachinidis A. Epigallocatechin 3 O Gallate inhibits the Ang II induced adhesion molecule expression in HUVEC via inhibition of MAPK pathways. *Cellular Physiology and Biochemistry*. 2007;20:859-66.
 21. Steffen Y, Schewe T, Sies H. (-)-Epicatechin elevates nitric oxide in endothelial cells via inhibition of NADPH oxidase. *Biochemical and biophysical research communications*. 2007;359(3):828-33.
 22. Steffen Y, Gruber C, Schewe T, Sies H. Mono-O-methylated flavanols and other flavonoids as inhibitors of endothelial NADPH oxidase. *Archives of biochemistry and biophysics*. 2008;469(2):209-19.
 23. Tsuneki H, Tokai E, Suzuki T, Seki T, Okubo K, Wada T, et al. Protective effects of coenzyme Q(10) against angiotensin II-induced oxidative stress in human umbilical vein endothelial cells. *European journal of pharmacology*. 2013.
 24. Unger RE, Krump-Konvalinkova V, Peters K, Kirkpatrick CJ. In Vitro Expression of the Endothelial Phenotype: Comparative Study of Primary Isolated Cells and Cell Lines, Including the Novel Cell Line HPMEC-ST1.6R. *Microvascular Research*. 2002;64(3):384-97.
 25. Mullen W, Edwards CA, Crozier A. Absorption, excretion and metabolite profiling of methyl-, glucuronyl-, glucosyl- and sulfo-conjugates of quercetin in human plasma and urine after ingestion of onions. *British Journal of Nutrition*. 2007;96(01):107.
 26. Wiczowski W, Romaszko J, Bucinski A, Szawara-Nowak D, Honke J, Zielinski H, et al. Quercetin from shallots (*Allium cepa* L. var. *aggregatum*) is more bioavailable than its glucosides. *The Journal of nutrition*. 2008;138:885-8.
 27. Wrangler K, Hermans JJR, Schiffers P, Moens AL, Paul M, Schmidt HHHW. Nox 1, 2, 4, 5: counting out oxidative stress. *British journal of pharmacology*. 2011;164(3).
 28. Davalos A, de la Pena G, Sanchez-Martin CC, Teresa Guerra M, Bartolome B, Lasuncion MA. Effects of red grape juice polyphenols in NADPH oxidase subunit expression in human neutrophils and mononuclear blood cells. *The British journal of nutrition*. 2009;102(8):1125-35.
 29. Appeldoorn MM, Venema DP, Peters TH, Koenen ME, Arts IC, Vincken JP, et al. Some phenolic compounds increase the nitric oxide level in endothelial cells in vitro. *Journal of agricultural and food chemistry*. 2009;57(17):7693-9.
 30. Vandenberg LN, Colborn T, Hayes TB, Heindel JJ, Jacobs DR, Jr., Lee DH, et al. Hormones and endocrine-disrupting chemicals: low-dose effects and nonmonotonic dose responses. *Endocr Rev*. 2012;33(3):378-455.
 31. Tribolo S, Lodi F, Winterbone MS, Saha S, Needs PW, Hughes DA, et al. Human metabolic transformation of quercetin blocks its capacity to decrease endothelial nitric oxide synthase (eNOS) expression and endothelin-1 secretion by human endothelial cells. *Journal of agricultural and food chemistry*. 2013;61(36):8589-96.
 32. Galindo P, Rodriguez-Gomez I, Gonzalez-Manzano S, Duenas M, Jimenez R, Menendez C, et al. Glucuronidated quercetin lowers blood pressure in spontaneously hypertensive rats via deconjugation. *PloS one*. 2012;7(3):e32673.
 33. Bakhshab S, Lari S, Ahmed F, Schulten H-J, Jayapal M, Karim S, et al. Endogenous controls in human umbilical vein endothelial cells under metabolic and oxidative stress. *BMC Genomics*. 2014;15(Suppl 2):P23.

Figure legends**Figure 1: Angiotensin II reduces NO[•] availability in HUVEC cultures.**

HUVEC cultures were incubated with Ang II for 1 – 24 h at A) 10 nM, B) 100 nM, C) 1000 nM and D) 10 000 nM, with NO[•] availability assessed post incubation in live cells using diaminofluorescein as described in the methods. E) HUVEC cultures were incubated with Ang II (100 nM, 8 h) ± the Angiotensin receptor type 1 antagonist Irbesartan (10 µM, 8 h). Filled bars indicate Ang II treatment (100 nM, 8 h). Graphs show mean values ± 1 S.D., n ≥ 3 independent experiments, * p<0.05.

Figure 2: Physiologically-obtainable quercetin concentrations prevent Ang II-induced decreases in nitric oxide production.

A) Ang II (100 nM, 8 h) induced decreases in NO[•] availability were restored to control levels by quercetin treatment (3 µM, 8 h), whereas quercetin treatment alone had no effect on NO[•] availability. This trend was observed for a range of physiologically-obtainable quercetin concentrations (3 nM – 3000 nM, B). C) endothelial nitric oxide synthase gene expression was not altered by either Ang II (100 nM, 8 h) or quercetin (3 µM, 8 h) treatments. Filled bars indicate Ang II treatment (100 nM, 8 h). Graphs show mean values ± 1 S.D., n ≥ 3 independent experiments, * p<0.05; ** p<0.01; *** p<0.001.

Figure 3: Quercetin prevents Ang II-induced superoxide production by NADPH oxidases.

A) Ang II-induced increases in intracellular superoxide are restored to control levels by quercetin co-treatment (8 h) at both 3 nM and 3 μ M concentrations, with no effect in quercetin only treated cells. B) Ang II-induced increases in intracellular superoxide are ameliorated to control levels by the NADPH oxidase inhibitor VAS-2870 (1 μ M, 8 h). Filled bars indicate Ang II treatment (100 nM, 8 h). Graphs show mean values \pm 1 S.D., $n \geq 3$ independent experiments, * $p < 0.05$.

Figure 4: Quercetin prevents Ang II-induced p47^{phox} expression at the protein and mRNA levels.

Quercetin co-treatment (3 μ M, 8 h) restored Ang II-induced p47^{phox} protein expression (100 nM, 8 h) to control levels in HUVEC cultures as assessed by Western blotting (A). This trend was reflected at the mRNA level, assessed using qPCR (B). The NADPH oxidase subunit gp91^{phox} was unaffected by either Ang II (100 nM, 8 h) or quercetin (3 μ M, 8 h) treatments (C). Filled bars indicate Ang II treatment (100 nM, 8 h). For the Western blotting data shown, the densitometry of the protein band of interest has been normalised to that of beta-actin, and it is this ratio that is presented in the bar graphs. Graphs show mean values \pm 1 S.D., $n \geq 3$ independent experiments, * $p < 0.05$; ** $p < 0.01$.

Ang II-induced PKC activity is not modulated by quercetin co-treatment

There was a trend of increase in PKC activity induced by Ang II exposure (100 nM, 8 h) but this effect was not ablated by quercetin co-treatment (3 μ M, 8 h). Filled bars indicate Ang II treatment (100 nM, 8 h). For this Western blotting data the densitometry values for each lane

of the PKC substrates blot has been normalised to that of tubulin. It is this ratio that is presented in the bar graphs. Graphs show mean values \pm 1 S.D., $n \geq 3$ independent experiments

Figure 6: Mechanistic insights into the effects of quercetin upon endothelial cell hypertension.

Quercetin restores angiotensin II-induced redox imbalance, centred around nitric oxide and superoxide. Changes in p47^{phox} protein expression are also restored to control levels, via modulation of gene expression, by a mechanism that seems independent of PKC activity. Arrows indicate the effects of angiotensin II treatment, which were ameliorated by quercetin co-exposure.


Supplementary Figure 1

A representative chromatogram of dihydroethidium (HE, precursor), the superoxide-specific reaction product of HE (2HE⁺), and fluorescein (internal standard). Dihydroethidium was detected at $m/z = 316.3$, 2HE⁺ at $m/z = 330$, and the internal standard fluorescein at $m/z = 333$.

Supplementary Figure 2

A) Cell media composition affects HUVEC response to Ang II (100 nM, 8 h). 24 h serum deprivation in M199 containing 0.5% (v/v) FBS resulted in a consistent decrease in NO[•] availability ($15 \pm 4\%$ of control) when HUVEC were treated with Ang II (100 nM, 8 h). Filled bars indicate Ang II (100 nM, 8 h) treatment. B) The effect of cell passage number upon Ang II (100 nM, 8 h) induced decreases in NO[•] availability. Passages 4-6 are demonstrated to be

599 responsive to Ang II, measured by NO[·] availability. C) The effect of confluency upon Ang II
 600 induced decreases in NO[·] availability. Cells grown for 2-3 days post seeding are demonstrated
 601 to be responsive to Ang II, measured by NO[·] availability, with decreasing responsiveness by 4
 602 days post seeding. D) The effects of DMSO and DMF upon Ang II induced decreases in NO[·]
 603 availability. It was shown that DMSO (0.1% v/v), but not DMF (0.1% v/v), prevented Ang II
 604 induced decreases in NO[·] availability. Thus DMF was used for all experiments. All
 605 experiments minimum n=3 independent experiments with mean \pm SD plotted. Filled bars
 606 indicates Ang II treatment. ** p<0.01.



Quercetin restores key cellular signalling molecules (nitric oxide and superoxide) to normal levels in a model of vascular endothelial cell (the cells that line blood vessels) dysfunction. This is achieved by restoring the levels of p47^{phox} (a key component in the superoxide generating machinery NADPH oxidase) to normal levels. The assessment of the modulation of these pathways by quercetin concentrations similar to those measured in human blood is a major part of the novelty of this work.

List of Abbreviations

Ang II – Angiotensin II

cDNA – Copy deoxyribose nucleic acid

DAF-2 DA – Diaminofluorescein diacetate

ddH₂O – distilled water

DHE - Dihydroethidium

DMF – Dimethyl-formamide

DMSO – Dimethyl-sulfoxide

DTPA – Diethylene triamine penta-acetic acid

ECGM – Endothelial cell growth media

ECL – Enhanced luminol chemiluminescence

eNOS – Endothelial nitric oxide synthase

FBS – Fetal bovine serum

HBSS – Hank's balanced salt solution

HRP – Horseradish peroxidase

HUVEC – Human umbilical vein endothelial cells

kDa – Kilo Dalton

LC-MS – Liquid chromatography mass spectrometry

L-NAME – NG-Nitro-L-arginine methyl ester

M199 – Medium 199

NADPH – Nicotinamide adenine dinucleotide 2'-phosphate (reduced form)

NO[•] - Nitric oxide

O₂^{•-} - Superoxide

PBS – Phosphate buffered saline

PKC – Protein kinase C






PVDF – Polyvinylidene difluoride


qPCR – Quantitative polymerase chain reaction




RNA – Ribose nucleic acid





ROS – Reactive oxygen species



SOD – Superoxide dismutase




















B-tubulin



Ang II	-	+	+	-
Quercetin	-	-	+	+